



XCI Encuentro anual de la Sociedad de Matemática de Chile

18 al 21 de Diciembre de 2023.

Departamento de Matemáticas, Facultad de Ciencias, Universidad de Chile.

Santiago, Chile.

Counting spanning subgraphs in dense hypergraphs

Matías Pavez-Signé*

Centro de Modelamiento Matemático (CNRS IRL2807)
Universidad de Chile

Abstract

We give a simple method to estimate the number of distinct copies of some classes of spanning subgraphs in hypergraphs with high minimum degree. In particular, for each $k \geq 2$ and $1 \leq \ell \leq k - 1$, we show that every k -graph on n vertices with minimum codegree at least

$$\begin{cases} \left(\frac{1}{2} + o(1)\right)n & \text{if } (k - \ell) \mid k, \\ \left(\frac{1}{\lceil \frac{k}{k-\ell} \rceil (k - \ell)} + o(1)\right)n & \text{if } (k - \ell) \nmid k, \end{cases}$$

contains $\exp(n \log n - \Theta(n))$ Hamilton ℓ -cycles as long as $(k - \ell) \mid n$. When $(k - \ell) \mid k$ this gives a simple proof of a result of Glock, Gould, Joos, Kühn and Osthus, while, when $(k - \ell) \nmid k$ this gives a weaker count than that given by Ferber, Hardiman and Mond or, when $\ell < k/2$, by Ferber, Krivelevich and Sudakov, but one that holds for an asymptotically optimal minimum codegree bound.

Trabajo realizado junto a:
Richard Montgomery¹

*Partially funded by Centro de Modelamiento Matemático (CMM) BASAL fund FB210005 for center of excellence from ANID-Chile and by the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (grant agreement No. 947978), e-mail: mpavez@dim.uchile.cl

¹Partially funded by the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (grant agreement No. 947978) and the Leverhulme Trust. Mathematics Institute, University of Warwick, UK